



<b>Title</b>	Infection Control Manual - MRSA Policy for NHS Borders
<b>Document Type</b>	Policy
<b>Document Number</b>	ICT001/08
<b>Version Number</b>	4 <sup>th</sup> Edition
<b>Approved by</b>	Infection Control Committee
<b>Issue date</b>	June 2014
<b>Review date</b>	June 2016
<b>Distribution</b>	All NHS Borders Staff
<b>Prepared by</b>	Infection Prevention Control Team
<b>Developed by</b>	Infection Prevention Control Team
<b>Equality &amp; Diversity Impact Assessed</b>	No

Item	Subject	Page
<b>1</b>	<b>Carriage, Infection and clearance</b>	3
1.1	Carriage	3
1.2	Infection	3
1.3	Clearance	3
<b>2.</b>	<b>Admission screening process</b>	4
2.1	Screening process by patient group	4
2.2	The clinical risk assessment (CRA)	4
2.3	MRSA screening samples	5
2.4	MRSA screening in the pre-admission unit	5
2.5	Management of colonised patients undergoing elective surgery	5
<b>3</b>	<b>Staff screening</b>	7
<b>4</b>	<b>Management of MRSA colonised patients in inpatient areas in BGH and Community Hospitals (excluding mental health)</b>	8
4.1	Hand hygiene	8
4.2	Placement	8
4.3	Contact screening of newly identified cases of MRSA colonization	8
4.4	Decolonisation of the MRSA positive patient	8
4.5	Informing a patient that they are colonised with MRSA	9
4.6	Transfer to another ward within the hospital	9
4.7	Personal protective equipment	9
4.8	Visitors	10
4.9	Crockery and cutlery	10
4.10	Linen	10
4.11	Waste	10
4.12	Equipment	10
4.13	Fans	10
4.14.	Slings	10
4.15	Other equipment	10
4.16	Routine cleaning	11
4.17	Terminal cleaning	11
<b>5</b>	<b>Transfer, discharge and movement of MRSA colonised and infected patients to theatre and diagnostic areas</b>	12
5.1	Diagnostic investigations	12
5.2	Theatre (including DPU)	13
5.3	Discharge of MRSA colonised patients	13
5.4	Outpatients	14
<b>6.</b>	<b>MRSA decolonisation treatment</b>	15
6.1	Topical decolonisation treatment	15
6.2	Post decolonisation screening	15

# 1. Carriage, Infection and clearance

## 1.1 Carriage

MRSA is carried at the same sites as meticillin sensitive *Staphylococcus*. (MSSA) The nose is the most common site of carriage with perineum, groin, axillae and throat being other common sites

*Staphylococcus aureus* including MRSA is also likely to be carried on areas of inflamed skin such as eczema / dermatitis and any wounds such as leg ulcers and pressure sores.

In addition the presence of a medical device which breaches the normal body defences such as peripheral venous cannulae or urinary catheters, will predispose to *Staphylococcus aureus* carriage at that site.

## 1.2 Infection

As distinct from carriage infection implies an invasive process to a greater or lesser degree and some degree of tissue inflammation. The common infections caused by *Staphylococcus aureus* including MRSA are skin infections such as boils and impetigo, cellulitis, osteomyelitis and infective endocarditis. In the healthcare setting *Staphylococcus aurea* often causes infection related to medical devices such as IV lines.

## 1.3 Clearance

*Staphylococcus aureus* carriage is normal for many people but in healthcare it can be useful to attempt to clear carriage with topical antimicrobials. (Decolonisation therapy). Sometimes people who were carriers of MRSA may lose carriage spontaneously, especially where the only site of carriage was a wound that has healed, or following decolonisation therapy. For the purposes of infection control management previously colonised patients can be regarded as clear if they have three clear screens from all appropriate sites taken at least one week apart and while not receiving topical or systemic antibiotic therapy that would suppress the growth in culture.

## 2. Admission Screening Process

MRSA screening consists of:

- The identification of patients at higher risk of MRSA colonisation by clinical risk assessment (CRA)
- The use of swabs from sites of usual MRSA carriage to detect carriage

### 2.1 Screening process by patient group

Patient group	Process
All adult emergency admissions except maternity and mental health	1. CRA 2. MRSA screening samples
All adult elective admissions with at least overnight stay anticipated	1. CRA 2. MRSA screening samples
All admissions to ward 9	If not performed for the admission in pre-assessment 1. CRA 2. MRSA screening samples
All admissions and transfers to ITU	1. CRA 2. MRSA screening samples
Patients transferred to paediatrics from another hospital	MRSA screening samples
Patients transferred to maternity from another hospital	MRSA screening samples
Paediatric admissions other than transfers from another hospital	Not screened
Admissions to mental health	Not screened
Admissions to community hospitals	Not screened

### 2.2 The clinical risk assessment (CRA)

Obtain answers to the 3 CRA questions

1. Has the patient previously been identified as MRSA positive/ (check clinical alert sheet on notes and flag on TrakCare)
2. Is the patient currently resident in a care home, institutional setting or transferred from another hospital?
3. Does the patient have a wound or device present e.g. Leg Ulcer, Pressure Sore, Hickman Line, PVC, urinary catheter?

If the answer is yes to any of these questions assume the patient to be MRSA positive and manage accordingly pending the results of MRSA screening samples.

### **2.3 MRSA screening samples**

These are taken from all patients admitted to acute adult wards except maternity, and all paediatric and adult transfers from other hospitals

Sites to be included in and MRSA screen

Nose swab (both anterior nares sampled using one swab)

Perineum swab

Throat swab (if patient will not accept perineal sampling or this is impractical for other reasons).

Wound swab(s)

Urine if patient is catheterized when admitted.

Sputum if the patient is expectorating.

### **2.4 MRSA screening in pre-admission unit**

Patients attending the pre-admission clinic for procedures that will require at least an overnight stay will be screened using the CRA and MRSA screening samples.

If a patient is negative for MRSA from their original pre-admission screen, and there is still a full negative response from the CRA questions including no previous history of MRSA, then this will be applicable for 18 weeks

If a patient has screened negative for MRSA at pre-assessment, but is positive for any of the CRA questions, then this will only be applicable for 8 weeks. Any patient with a positive response to their CRA question are assumed to be more at risk of acquisition

### **2.5 Management of colonised patients undergoing elective surgery**

Colonised patients undergoing elective surgery should be offered decolonization therapy applied either pre-operatively or peri-operatively. Peri-operative decolonization should be commenced two or three days prior to the date of surgery in order to effectively suppress microbial load and reduce the risk of infection.

If antimicrobial prophylaxis is needed for the procedure this should be adjusted to include cover for MRSA as specified in the NHS Borders Antimicrobial Guidelines for hospitals available at:  
<http://intranet/resource.asp?uid=2845>

### 3. Staff screening

This is very seldom necessary but may be useful when unexplained acquisition of MRSA occurs within NHS Borders patients and personnel. The decision to screen will be agreed with senior medical and nursing staff within the involved clinical board. The process will be co-ordinated by Occupational Health and the Infection Prevention Control Team. Staff found to be MRSA positive will be seen and counselled by a member of Occupational Health staff. The appropriate decolonisation treatment will be prescribed and provided, and follow up screening organised.

Occupational Health will advise when it is appropriate to return to work.

## **4. Management of MRSA colonised patients in inpatient areas in BGH and Community Hospitals (excluding mental health)**

### **General Measures**

#### **4.1 Hand hygiene**

Hand hygiene (either hand washing or application of alcohol gel) is the single most important measure for prevention of transmission of MRSA in clinical settings.

Please refer to [Standard Infection Control Precautions](#) and [NHS Borders Zero Tolerance Hand Hygiene Policy](#).

#### **4.2 Placement**

Patients colonized with MRSA should be nursed in a single room unless risk assessment shows that this compromises patient care.. The room should be identified as one being used for isolation and the door remain closed unless risk assessment shows that this compromises patient care. Any such assessment should be documented in the unitary record

#### **4.3 Contact screening of newly identified cases of MRSA colonization**

Identification of MRSA colonisation or infection in patients some time following admission will often imply transmission within the ward. It may then be appropriate to screen contacts of the index case to prevent further transmission within the ward. When new cases such as these are identified the need for contact screening will be considered by the IPCT in discussion with ward staff.

#### **4.4 Decolonisation of the MRSA positive patient**

The usual approach should be to use topical decolonization on patients colonized with MRSA who are admitted to hospital as this appears to reduce the risk of MRSA infection during their admission. There may be factors such as skin sensitivity or multiple wounds which would make it less applicable in a particular patient. The Infection Prevention Control Team will advise you if an attempt should be made to decolonise patients known to be positive with



MRSA. Please remember that, as with any other procedure in hospital, any proposed investigation or treatment should be adequately explained and discussed with the patient and/ or their relatives, and that they have the right to decline such intervention.

#### **4.5 Informing a patient that they are colonized with MRSA**

In the first instance this should be undertaken by a member of the Medical/ Nursing staff caring for the patient. However, after this the IPCT are happy to speak to patients/ relative if this is deemed appropriate or further questions/ help is required.

Relatives should only be informed with the agreement/ knowledge of the patient. For children, the parents/ guardian will be informed. Colonised patients and where appropriate their relatives/carers should be offered written information on MRSA.

#### **4.6 Transfer to another ward within the hospital**

Transfer of MRSA affected patients to other wards should be minimised to reduce the risk of spread, but this should not compromise other aspects of the patient's care, such as rehabilitation. In all cases, it is the responsibility of ward and clinical staff to inform relevant departments of a patient's MRSA status well in advance of transfer

### **Specific measures**

#### **4.7 Personal protective equipment (PPE)**

Glove and apron (disposable)

Must be worn, by all staff, for all procedures that involve patient contact. Gloves and apron must be removed before leaving the patient surroundings. Wearing gloves does not preclude the need for hand washing.

Masks and eye protection

Should be worn if procedures are to be undertaken in which there is judged to be a significant risk of splashing of blood/body fluids or when the patient is unable to control cough and sputum is colonized.

#### **4.8 Visitors**

There is no need for visitors to wear PPE unless undertaking patient care. They should, however, be encouraged to clean their hands on leaving the patient's surroundings.

#### **4.9 Crockery and cutlery**

Use normal utensils. Wash in dishwasher.

#### **4.10 Linen**

Treat as infected. For personal laundry follow any local arrangements in place. Change linen and clothing on a daily basis.

#### **4.11 Waste**

Should be treated as clinical waste i.e. placed in a yellow clinical waste bag.

#### **4.12 Equipment**

Clean with general-purpose detergent (GPD) and warm water. For equipment to be removed from room, clean with 1,000ppm solution.

#### **4.13 Fans**

Portable fans should not be used close to a patient known to be MRSA positive when nursed in a bay with other patients.

#### **4.14 Slings**

Single patient use (disposable) slings should be used.

#### **4.15 Other equipment**

Non-critical items of healthcare equipment in direct contact with the patient (e.g. stethoscopes, BP cuffs) should, where possible, be dedicated for that patient during their hospital stay. Such items should

be appropriately decontaminated or disposed of after the patient is discharged

## **Cleaning**

### **4.16 Routine cleaning**

Routine cleaning of siderooms occupied by colonized patients should be with detergent.

### **4.17 Terminal cleaning**

On transfer/discharge the patient's furniture should be cleaned with 1,000ppm Chlorine solution. The floors should also be washed with 1,000ppm Chlorine solution. Ensure laundering of curtains.

## 5 Transfer, discharge and movement of MRSA colonised and infected patients to theatre, diagnostic areas and outpatients.

MRSA infection/ colonization should not interfere with the management of the patient. When visiting a diagnostic or therapeutic department, the department must be informed, in advance, so that appropriate infection control measures for that department can be implemented.

### 5.1 Diagnostic Investigations

Occlude any lesions whenever possible with an impermeable dressing. Attendants who may be in contact with the patient should wear disposable plastic aprons to protect their clothing. Aprons should be removed when contact with the patient has finished and disposed of as clinical waste. Gloves need only be worn if staff transporting the patient is specifically instructed to do so by the nurse in charge, or by the Infection Prevention & Control Team.

Linen should be treated as used and disposed of in accordance with NHSB Laundry Policy.

Chairs/trolleys should be wiped down with general purpose detergent and water (or detergent wipes) after transfer or use. If a patient is being transferred on their bed, both bed and linen should be clean prior to transfer.

Patient should spend the minimum time in the department, being sent for when the department is ready and not left in a waiting area with other patients.

Staff coming into direct contact with the patient should wear a disposable plastic apron and gloves. Staff should avoid direct contact with other patients whilst dealing with an MRSA positive patient.

Equipment and the number of staff attending should be kept to a minimum.

Surfaces with which the patient has had direct contact should be wiped clean with a general purpose detergent and water.

Linen should be disposed of in accordance with the NHSB Laundry Policy

## 5.2 Theatre (including DPU)

Occlude any lesions whenever possible with an impermeable dressing.

Attendants who may be in contact with the patient should wear disposable plastic aprons to protect their clothing. Aprons should be removed when contact with the patient has finished and disposed of as clinical waste. Gloves need only be worn if staff transporting the patient is specifically instructed to do so by the nurse in charge, or by the Infection Prevention & Control Team.

Chairs/trolleys should be wiped down with general purpose detergent and water (or detergent wipes) after transfer or use. If a patient is being transferred on their bed, both bed and linen should be clean prior to transfer.

Patient should spend the minimum time in the department, being sent for when the department is ready and not left in a waiting area with other patients.

Staff coming into direct contact with the patient should wear a disposable plastic apron and gloves. Staff should avoid direct contact with other patients whilst dealing with an MRSA positive patient.

Equipment and the number of staff attending should be kept to a minimum.

Surfaces with which the patient has had direct contact should be wiped clean with a general purpose detergent and water.

Linen should be disposed of in accordance with the NHSB Laundry Policy

MRSA colonised patients do not need to be placed at the end of a theatre list and can be recovered in the main recovery areas.

## 5.3 Discharge of MRSA colonised patients

The General Practitioner and other health care agencies involved in the patient's care should be informed of new recognition of MRSA colonisation. The ward nurses should inform community nurses where ongoing care is required

MRSA carriers will not normally require special treatment after discharge from hospital. Advice may be obtained from the Infection

Prevention & Control Team if patients due for transfer or discharge are undergoing topical MRSA eradication therapy.

Patients and their carers should be fully informed about their MRSA status prior to discharge and reassured that their healthy relatives and contacts are not at risk. Advice may be sought from the Infection Prevention & Control Team if patients or their carers identify contacts that may be at risk due to their own health, e.g. contacts with lowered immunity or chronic skin lesions

Patients should be advised that they should inform staff at any future hospital admission that they have previously been identified as carriers of MRSA.

Patients who have died colonised or infected with MRSA do not require to be placed in body bags for this reason as they do not pose a risk to mortuary staff, patients' relatives, or undertakers.

#### **5.4 Outpatients**

Colonised patients attending out-patient appointments should attend as normal and not segregated. Gloves and aprons should be worn if performing a clinical examination.

## 6. MRSA DECOLONISATION TREATMENT

Decolonisation treatment will last 5 days. It will normally be applied to inpatients at BGH found to be colonized with MRSA, usually at the direction of the IPCT, but may also be initiated by the inpatient team. Normally a maximum of only two consecutive treatments will be given in a particular admission, but contact IPCT for advice.

### 6.1 Topical decolonisation treatment

1. Apply Mupirocin (Bactroban) nasal ointment three times a day to the inner surface of the nostrils (use a cotton wool bud to enhance application). If the patient's MRSA strain is resistant to mupirocin, discuss with the IPCT for alternative treatment
2. Apply daily 4% Chlorhexidine cleansing solution instead of soap. Rinse off and towel dry
3. Chlorhexidine cleanser should also be used at least twice weekly as a shampoo whilst on the decolonisation treatment (hair conditioner may be used for the final rinse)
4. Alternative treatments are available for patients whose skin cannot tolerate chlorhexidine, for further information contact the IPCT
5. The patient should use chlorhexidine 0.2% mouthwash two times daily

The patient's bed linen (and night wear if possible) should be changed daily during the decolonization treatment.

Alternative products are available for patients with chlorhexidine allergy or sensitivity. Contact the IPCT for advice if needed.

### 6.2 Post-decolonisation screening

- This should be commenced no sooner than 48 hours after the treatment regime has been completed
- If patient commences or remains on antibiotic treatment active against MRSA, delay screening until 48 hours after antibiotics have been discontinued

- Mark specimen 'MRSA clearance screening'
- If the first screen is negative, consult the Infection Prevention Control Team who will reassess the need for further screening
- If the first screen is positive consult the Infection Prevention Control Team who will reassess the need for further treatment.

**A negative screen following decolonization does not guarantee that the patient will not recolonise at a later date. The patient record/ notes continue to alert of the history of MRSA carriage**